

FITNESS® Fungicide will provide early control of diseases on a wide range of crops without causing injury. FITNESS Fungicide helps maintain a healthy flag leaf, protecting from leaf diseases and protecting your investment.

CROPS:

• Wheat, barley, oats, corn, canola, soybeans (grown for seed), dry bean*, canaryseed, timothy

*Control of rust (Uromyces spp.)

FEATURES:

- Wide spectrum of registered crops
- Wide spectrum of registered diseases
- Systemic activity

WATER VOLUMES:

Ground: 20 gal/ac (200 L/ha) **Aerial:** 5 gal/ac (50 L/ha)

APPLICATION RATES:

MIXING ORDER:

- 1. Fill tank half full with clean water. Engage gentle agitation.
- 2. Add the required amount of Fitness and agitate thoroughly.
- 3. Continue filling the tank with water until it is ninetenths full and, if applicable, add tank-mix partner.
- 4. Complete filling the tank with water, maintaining agitation throughout mixing and spraying.

TANK MIXES:

- Logic M, Brotex 240, Pardner, 2,4-D Amine, 2,4-D Ester, MCPA Amine, MCPA Ester, Foothills NG (wheat only), Horizon NG (wheat only), Axial or Buctril M.
- Fitness may be applied with up to 9 lb/ac (4 kg/ac) of actual nitrogen. The appropriate amount of urea can be dissolved in water and added to the spray tank before adding the fungicide.

CAUTION: Excessive nitrogen concentrations may injure the crop.

NOTE: Do not add nitrogen when tank-mixing Fitness fungicide with a herbicide.

• Do not tank mix herbicides for application on oats.

lutrien

a Solutions

ACTIVE INGREDIENT:

Propiconazole 418 g/L - Group 3

PACK SIZES:

4.8 L jug (4 jugs per case)

CROP	DISEASES	RATE (ml/ac)	TIMING
Wheat (spring, winter, durum)	For suppression of: Septoria leaf blotch (<i>Septoria tritici</i>) Tan Spot (<i>Pyrenophora tricitici-repentis</i>)	60-120	 Early application: The first sign of disease, usually at the beginning of stem elongation. Later application: Before head is half emerged.
	For control of: Septoria leaf blotch <i>(Septoria tritici)</i>	120	
Barley	Net blotch (Drechsiera teres)	120	
Oats	Septoria leaf blotch (Septoria tritici)	120	
Canola	Blackleg (Leptosphaeria macufans)	120	Apply during the rosette stage. Between second true leaf and bolting.

*For more crops, diseases, rates and timing, refer to product label



ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

© 2020 Loveland Products, Inc. All Rights Reserved. All of the trademarks and service marks displayed are marks of their respective owners.

