

GROUP 3 FUNGICIDE

Propel[®]

FUNGICIDE

COMMERCIAL

Emulsifiable Concentrate for:
Broad-spectrum disease control in Wheat, Barley, Oats, Canola, Corn, Legume Vegetables including Soybeans and for the suppression of Septoria Leaf Mottle on Canaryseed.

FOR SALE FOR USE ON TIMOTHY HAY IN THE PRAIRIE PROVINCES ONLY.

GUARANTEE:

Propiconazole250 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING



WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

REGISTRATION NO.: **29548**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L to 1000 L**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON
N1G 4Z3
Telephone: 1-877-964-3682

Label

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IF POISONING IS SUSPECTED, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote for this product. Apply symptomatic therapy. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

PRECAUTIONS

KEEP OUT OF THE REACH OF CHILDREN. DO NOT get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin sensitization reactions in certain individuals.

Wear long pants, a long sleeve shirt, shoes and socks and chemical-resistant gloves during mixing/loading, application, clean-up and repair activities. Wear protective goggles or face shield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used. **DO NOT** eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes. After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse.

DO NOT contaminate food or feed.

DO NOT allow entry into treated area for 12 hours following application (12 hr REI). See the **DIRECTIONS FOR USE** section for crop-specific restricted entry intervals.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. This pesticide is toxic to fish. DO NOT spray any body of water by direct application, drift or by cleaning and rinsing spray equipment. This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning of equipment.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application sites such as hedgerows and woodland.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL OR REFILLING:

For recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Dispose of the rinsings in accordance with provincial requirements.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

DO NOT reuse this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

For returnable containers:

DO NOT reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

It must be noted Propel[®] is a trademark of Viterra. Viterra is a licensed user.

GROUP 3 FUNGICIDE

Propel[®]

FUNGICIDE

COMMERCIAL

Emulsifiable Concentrate for:

Broad-spectrum disease control in Wheat, Barley, Oats, Canola, Corn, Legume Vegetables including Soybeans and for the suppression of Septoria Leaf Mottle on Canaryseed.

FOR SALE FOR USE ON TIMOTHY HAY IN THE PRAIRIE PROVINCES ONLY.

GUARANTEE:

Propiconazole250 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

CAUTION



POISON

WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

REGISTRATION NO.: **29548**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON
N1G 4Z3
Telephone: 1-877-964-3682

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IF POISONING IS SUSPECTED, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote for this product. Apply symptomatic therapy. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

PRECAUTIONS

KEEP OUT OF THE REACH OF CHILDREN. DO NOT get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin sensitization reactions in certain individuals.

Wear long pants, a long sleeve shirt, shoes and socks and chemical-resistant gloves during mixing/loading, application, clean-up and repair activities. Wear protective goggles or face shield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used. **DO NOT** eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes. After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse.

DO NOT contaminate food or feed.

DO NOT allow entry into treated area for 12 hours following application (12 hr REI). See the **DIRECTIONS FOR USE** section for crop-specific restricted entry intervals.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. This pesticide is toxic to fish. DO NOT spray any body of water by direct application, drift or by cleaning and rinsing spray equipment. This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning of equipment.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application sites such as hedgerows and woodland.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL OR REFILLING:

For recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Dispose of the rinsings in accordance with provincial requirements.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

DO NOT reuse this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

For returnable containers:

DO NOT reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

DIRECTIONS FOR USE

DO NOT use in greenhouses.

A restricted entry interval (REI) of 1 day is required for workers hand-harvesting and detasseling treated corn.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

GENERAL INFORMATION

DO NOT graze animals on treated green crops within three days of application of Propel® Fungicide.

PRODUCT INFORMATION

Propel Fungicide is a broad spectrum systemic fungicide for control of a wide range of diseases on certain crops. Propel Fungicide will protect the crop from yield and quality losses due to disease. Propel Fungicide may be used in conjunction with higher seeding rates, higher fertilizer inputs, plant growth regulators and other fungicides as required.

(See "NOTE" under PRECAUTIONS)

Crop Information:

Propel Fungicide may be applied to:

- Barley, Spring
- Canary Seed
- Canola
- Corn (including Field, Seed, and Sweet)
- Legume Vegetables
- Oats
- Soybeans
- Timothy
- Wheat, Spring and Winter (including Hard Red, Durum, Canada Prairie, Soft White)

Diseases Controlled:

Barley, Spring:	Net Blotch; Spot Blotch; Scald; Powdery Mildew; Septoria Leaf Spot; Leaf Rust; Stem Rust
Canary Seed:	Septoria Leaf Mottle
Canola:	Blackleg
Corn:	Rusts; Northern Corn Leaf Blight; Southern Corn Leaf Blight; Helminthosporium Leaf Spot; Eye Spot; Grey Leaf Spot
Legume Vegetables including Soybeans:	Rust (<i>Phakopsora</i> spp.); Cercospora Leaf Spot (<i>Cercospora</i> spp.); Powdery mildew (<i>Microsphaera</i> spp., <i>Erysiphe</i> spp.)
Oats:	Septoria Leaf Blotch; Crown Rust
Soybeans grown for seed:	Frogeye Leaf Spot (<i>Cercospora</i> spp.), Aerial Web Blight (<i>Rhizoctonia solani</i>)
Timothy:	Purple eyespot (<i>Cladosporium phlei</i>)
Wheat, Spring and Winter: (including Hard Red, Durum, Canada Prairie, Soft White)	Septoria Leaf Spot; Septoria Glume Blotch; Powdery Mildew; Leaf and Stem Rust; Tan Spot; Stripe Rust

Factors Affecting Propel Fungicide Performance:

Propel Fungicide should be applied as a preventative disease control measure. Established diseases are more difficult to control and may have already reduced crop vigour. If rainfall occurs within one hour of application, reapplication is necessary.

Field Sprayer Application:

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial Application:

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor span. High humidity and low temperatures (10 - 20°C) allow for a better deposition of spray droplets.

Sprayer and Application Information:

The performance of this product depends on correct application. Follow the guidelines given below for optimal application of Propel Fungicide.

Sprayer Information:

	GROUND APPLICATION	AERIAL APPLICATION
Spray Volume:	minimum 200 L water/ha	40 - 50 L water/ha
Spray Pressure:	200 - 300 kPa	100 - 200 kPa
Nozzle Type:	110° Flat Fan (XR11004, 4110-20)	Flat Fan (6510-6515) or Hollow Cone (D8-45)
Droplet Size:	Medium Spray (300 - 400 microns VMD)	Medium Spray (350 - 400 microns VMD)
Ground Speed:	10 km/h	
Nozzle Angle:	90° (straight down)	90° (straight down)
Boom Height:	40 - 50 cm above the crop canopy	2 - 3 m above the crop canopy

Ground Application - Mixing and Spraying Instructions

- 1) Spray equipment should be thoroughly flushed with clean water before mixing Propel Fungicide.
- 2) Fill spray tank 1/2 full with clean water. Engage gentle agitation.
- 3) Add the required amount of Propel Fungicide and agitate thoroughly.
- 4) Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tankmix partner.
- 5) Complete filling the spray tank with water, maintaining agitation during mixing and spraying operations.
- 6) Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Aerial Application - Mixing and Spraying Instructions

- 1) Spray equipment should be thoroughly flushed with clean water before mixing Propel Fungicide.
- 2) Fill premix tank 1/2 full with clean water. Engage gentle agitation.
- 3) Add the required amount of Propel Fungicide and agitate thoroughly.
- 4) Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tankmix partner.
- 5) Complete filling the premix tank with water.

- 6) Maintain gentle agitation during mixing.
- 7) Transfer the premix contents into the aircraft spray tank.
- 8) Maintain sufficient agitation during the mixing and spraying operation to ensure Propel Fungicide remains in suspension.
- 9) Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for a specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL APPLICATION USE PRECAUTIONS:

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift; therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as a shelter-belt) or aquatic habitat.

AERIAL APPLICATION OPERATOR PRECAUTIONS:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the label recommendations on the ground application section. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Buffer zones:

Use of the following spray methods or equipment. **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				Terrestrial habitat
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer*	Beans, peas, soybeans, chickpeas, corn, wheat, oats, canary seed, canola, barley, rye, triticale		1	0	1	1	1
Aerial	Beans, corn, oats, wheat, barley, triticale (seed prod.)	Fixed wing	1	0	3	1	20
		Rotary wing	1	0	1	1	20

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

For tank mixes, consult the labels of tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

PRODUCT SPECIFIC PRECAUTIONS:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-87-SYNGENTA (1-877-964-3682) or obtain technical advice from the distributor or your provincial agricultural representative.

















Fertilizer - Mixing and Spraying Instructions

If desired, small amounts of nitrogen may be applied with Propel Fungicide. The appropriate amount of urea can be dissolved in water and added to the spray tank before adding Propel Fungicide. The rate of actual nitrogen must not exceed 10 kg/ha.

CAUTION: Excessive nitrogen concentrations may injure the crop.

NOTE: DO NOT add nitrogen when tank-mixing Propel Fungicide with a herbicide.

**WHEAT, BARLEY, OATS
CEREAL GROWTH STAGES***

							
3 leaves unfolded 13	Main shoot and tiller 21	Main shoot and 4 tillers 24	Leaf sheath length to ligula 30	Pseudo stem anetion 30	1st node detectable 31	2nd node detectable 32	Flag leaf just visible 37
							
Ligule of last leaf visible 39	Boots swollen 45	Ear just visible 51	Ear 1/2 emerged 55	Ear fully emerged 59	Flowering over at base of ear 69	Flowering over (anthesis) watery ripe 71	Grain hard 81

WHEAT, BARLEY, OATS	
INSTRUCTIONS FOR USE:	
Apply Propel Fungicide at the very early stages of disease. This could occur anytime during tillering or stem elongation. Typically, an application from the beginning of stem elongation up to flag leaf emergence is required.	
Propel Fungicide lasts about three weeks in the plant. If conditions favourable to disease continue after this length of time, another application will be necessary to maintain control. The second spray is usually applied at the time of head emergence. In most cases, this second application is essential to maintain control of the <i>Septoria</i> disease complex.	
LAST APPLICATION MUST BE MADE PRIOR TO 45 DAYS BEFORE HARVEST (45 day PHI).	
CROP/DISEASES	<u>WHEAT: Septoria Leaf Spot Tan Spot</u> <u>SPRING BARLEY: Net Blotch</u>
RATE/HA	250 - 500 mL
EARLY Application	At G.S. 12-23 (as early as the two leaf stage). For early season disease suppression, use the lower rate for suppression under normal field conditions. Use the higher rate for control if there is a history of high disease pressures in the field and/or field conditions favour disease development.
LATER Application	At the first sign of disease (G.S. 29-37) or before head is half emerged (G.S. 49-55). Apply only the high rate on any application from G.S. 29-55.
MAXIMUM NUMBER OF APPLICATIONS PER SEASON	2
CROP/DISEASES	<u>WHEAT: Septoria Leaf Spot and Glume Blotch Powdery Mildew Leaf and Stem Rust Tan Spot Stripe Rust</u> <u>SPRING BARLEY: Net Blotch Spot Blotch Scald Powdery Mildew Septoria Leaf Spot Leaf and Stem Rust</u> <u>OATS: Septoria Leaf Blotch Crown Rust</u>
RATE/HA	500 mL
EARLY Application	At the first sign of disease, usually at the beginning of stem elongation (G.S. 29-37).
LATER Application	Before head is half emerged (G.S. 49-55).
MAXIMUM NUMBER OF APPLICATIONS PER SEASON	2

HERBICIDE TANK-MIXING - WHEAT & BARLEY:

Propel Fungicide can be tank-mixed with ONLY ONE of these herbicides:

2,4-D Amine MCPA Amine
 Buctril® M Pardner®
 HORIZON® 240EC Herbicide (Wheat Only)
 AXIAL® 100EC Herbicide (Spring Wheat and Spring Barley Only) (FOR USE ONLY IN THE PRAIRIE AND INTERIOR OF BRITISH COLUMBIA)

Tank-mixing Precautions:

1. DO NOT tank-mix Propel Fungicide with herbicides for application onto Oats.
2. Weeds and crops must be at the correct stage of growth as specified in both the Propel Fungicide label and the tank mix partner label.
3. 2,4-D and MCPA formulations may be applied either by ground application or aerial

application; tank-mixtures of Propel Fungicide and Buctril M, Pardner, or AXIAL 100EC Herbicide can only be applied by ground application.

4. For fields treated with AXIAL 100EC Herbicide, no crop may be seeded until the following year. There are no crop rotation limitations the year following application of Axial 100EC Herbicide.
5. Tank mixing with the herbicide AXIAL 100EC will slightly decrease Propel efficacy.
6. Consult the label of the herbicide partner for a list of weeds controlled, directions for use and precautions.
7. When tank mixing, adhere to the most restrictive label limitations and precautions.
8. Compatibility should always be confirmed by premixing small proportional quantities of water, Propel Fungicide, and the tank-mix partner in advance.

NOTE:

- DO NOT graze animals on treated green crops within three days of application of Propel Fungicide.
- DO NOT feed straw from crops treated with herbicide tank mixes to livestock.

HERBICIDE TANK-MIXING – WHEAT ONLY:

FOR USE ONLY IN THE PRAIRIE AND INTERIOR OF BRITISH COLUMBIA.

Propel Fungicide can be tank-mixed with HORIZON 240EC Herbicide for disease and grassy weed control.

Tank-mixing Precautions:

1. DO NOT apply by air.
2. Consult the label of HORIZON 240EC Herbicide for a list of weeds controlled, directions for use and precautions.
3. Apply prior to emergence of the 4th tiller (herbicide timing).

TIMOTHY	
<u>INSTRUCTIONS FOR USE:</u>	
Apply Propel at the very early stages of disease. If conditions favourable to disease continue after this length of time, another application will be necessary to maintain control. DO NOT APPLY BY AIR. LAST APPLICATION MUST BE MADE PRIOR TO 14 DAYS BEFORE HARVEST (14 day PHI).	
DISEASE	Purple eyespot
RATE/HA	500 mL
EARLY Application	At the first sign of disease, usually at the beginning of flowering (G.S. 59-61).
LATER Application	Full flowering (G.S. 65-73).
Minimum Interval between Applications	14 days
Maximum Seasonal Application Rate	2 applications at 500 mL/ha (1.0 L/ha in a season).

CANOLA				
INSTRUCTIONS FOR USE:				
Propel Fungicide will control blackleg and enhance yield potential during the early stages of canola growth. The disease may reappear later in the season, but with minimal effect on yield. LAST APPLICATION MUST BE MADE PRIOR TO 60 DAYS BEFORE HARVEST (60 day PHI).				
DISEASES	RATE/HA	REMARKS/TIMING		
Blackleg	500 mL	Apply during the rosette stage: between 2 nd true leaf and bolting.		
		Seedling	Rosette	Bud (Bolted)
		Stage 1	Stage 2	Stage 3

TANK-MIXING – SEED CORN, FIELD CORN and SWEET CORN:

Propel Fungicide can be tank-mixed with ONLY ONE of the following partners:

MATADOR[®] 120EC Insecticide

WARRIOR[®] Insecticide

Ripcord[®]

Tank-mixing Precautions:

1. DO NOT APPLY THE TANK MIX WITH WARRIOR INSECTICIDE BY AIR.
2. The tank mix of Propel Fungicide + MATADOR 120EC Insecticide or Propel Fungicide + Ripcord can be applied by air and ground. Use 40 L of water per hectare when applying by air.
3. Insects and crops must be at the correct stage as specified on the Propel Fungicide as well as MATADOR 120EC, WARRIOR Insecticide and Ripcord (PCP 15738) labels. Follow the directions for use and precautions on all labels.
4. DO NOT apply within 14 days of harvest (14 day PHI) when using these tank-mixes on field and sweet corn.
5. Compatibility should always be confirmed by premixing small proportional quantities of water, Propel Fungicide, and the tank-mix partner in advance.

TANK MIXES

SEED CORN	
DISEASES	Rusts
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when rust pustules first appear. Under severe disease pressure, make a second application 14 days after. Under severe disease pressure, make a third application 14 days after. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Northern Corn Leaf Blight Southern Corn Leaf Blight Helminthosporium Leaf Spot
RATE/HA Propel Fungicide alone	250 - 500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 250 - 500 mL/ha of Propel Fungicide when disease first appears. Use the 250 mL rate if disease pressure is low. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Eye Spot Grey Leaf Spot
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when disease first appears. DO NOT apply the tank mix with WARRIOR Insecticide by air.
FIELD CORN and SWEET CORN	
DISEASES	Rusts
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when rust pustules first appear. Under severe disease pressure, make a second application 14 days after. DO NOT apply the tank mix with WARRIOR Insecticide by air.

FIELD CORN and SWEET CORN	
DISEASES	Northern Corn Leaf Blight Southern Corn Leaf Blight Helminthosporium Leaf Spot
RATE/HA Propel Fungicide alone	250 - 500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 250 - 500 mL/ha of Propel Fungicide when disease first appears. Use the 250 mL rate if disease pressure is low. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Eye Spot Grey Leaf Spot
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when disease first appears. DO NOT apply the tank mix with WARRIOR Insecticide by air.

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS

Soybeans

Edible-podded legume vegetables - Any succulent cultivar of edible podded bean (*Phaseolus* spp.) and any succulent cultivar of edible-podded pea (*Pisum* spp.): Bean (*Phaseolus* spp.) (includes runner bean, snap bean, wax bean); bean (*Vigna* spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean); jackbean; pea (*Pisum* spp.) (includes dwarf pea, edible-podded pea, snow pea, sugar snap pea); pigeon pea; soybean (immature seed); and sword bean.

Succulent shelled pea and bean - Any succulent shelled cultivar of bean (*Phaseolus* spp.) and garden pea (*Pisum* spp.): Bean (*Phaseolus* spp.) (includes lima bean, green bean); broad bean (succulent); bean (*Vigna* spp.) (includes black-eyed pea, cowpea, southern pea); pea (*Pisum* spp.) (includes English pea, garden pea, green pea); and pigeon pea.

Dried shelled pea and bean - Any dried cultivar of bean (*Phaseolus* spp.) and dried cultivar of pea (*Pisum* spp.): Dried cultivars of bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and sweet white lupin); (*Vigna* spp.) (includes adzuki beans); (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean, mung bean, rice bean, southern bean, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; pea (*Pisum* spp.) (includes field pea); and pigeon pea.

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS	
DISEASE	Asian (Soybean) Rust (<i>Phakopsora pachyrhizi</i>)
CROPS	All Crop Group 6 legume vegetables listed above, and soybean
RATE/HA	500 – 756 mL
REMARKS	<p>For control of Asian Rust, apply the lower rate under conditions of low disease pressures.</p> <p>Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.</p> <p>APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).</p>
DISEASE	Powdery Mildew (<i>Microsphaera diffusa</i> , <i>Erysiphe pisi</i> , <i>E. polygoni</i>)
CROPS	All Crop Group 6 legume vegetables listed above, and soybean
RATE/HA	500 mL
REMARKS	<p>Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.</p> <p>APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).</p>
DISEASE	Cercospora Leaf Spot (<i>Cercospora kikuchi</i>)
CROPS	Soybeans only
RATE/HA	500 mL
REMARKS	<p>Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.</p> <p>APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).</p>

TANK MIX - Propel FUNGICIDE with QUADRIS® Flowable Fungicide

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS	
DISEASE	Asian (Soybean) Rust (<i>Phakopsora pachyrhizi</i>)
CROPS	All Crop Group 6A, B, C legume vegetables listed above, and soybeans
RATE/HA	0.5 - 0.75 L/ha of Propel Fungicide + 0.3 - 0.45 L/ha of QUADRIS Flowable Fungicide
REMARKS	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application at 14 days interval may be needed if conditions persist. It is important to protect the developing pod of soybean and podded legume vegetables. Good spray coverage and canopy penetration are important for best results. Apply in a minimum of 45 L of water per hectare.
Restrictions: DO NOT APPLY MORE THAN 2 APPLICATIONS OF Propel FUNGICIDE PER SEASON. DO NOT APPLY MORE THAN 2 APPLICATIONS OF QUADRIS FLOWABLE FUNGICIDE PER SEASON. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables). DO NOT make more than one application to soybean hay and dry pea hay. DO NOT apply within 14 days of harvest (14 day PHI) of soybean hay and dry pea hay. DO NOT feed dried pea vines to livestock. Not all members of the legume vegetable group have been tested for efficacy and phytotoxicity at the recommended label rates and this tank mix should be used at the discretion of the user. The tank mix of Propel Fungicide plus QUADRIS Flowable Fungicide may be applied by air or ground application equipment.	

TANK MIX - Propel FUNGICIDE with MATADOR 120EC Insecticide or WARRIOR Insecticide

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS
Propel Fungicide can be tank mixed with MATADOR 120EC Insecticide or WARRIOR Insecticide for foliar disease and insect control. Apply Propel Fungicide at a rate of 250 - 756 mL/ha in a tank mix with MATADOR 120EC Insecticide or WARRIOR Insecticide at a rate of 83 - 233 mL/ha for control of soybean aphid on soybean and at a rate of 83 mL/ha for the rest of the crops in Crop Group 6. Refer to both the Propel Fungicide, MATADOR 120EC Insecticide, or WARRIOR Insecticide labels for diseases and insects controlled, specific application instructions, and precautions. Pests and crops must be at the correct stage as specified on the Propel Fungicide, MATADOR 120EC Insecticide, or WARRIOR Insecticide labels. DO NOT apply more than 2 applications per season of this tank mix. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables). DO NOT graze or harvest treated forage, straw or hay for livestock feed. Not all members of the legume vegetable group have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user. This tank mix can be applied by ground application equipment only. DO NOT APPLY THE TANK MIX WITH MATADOR 120EC Insecticide or WARRIOR Insecticide BY AIR.

TANK MIXES - Propel FUNGICIDE with QUADRIS Flowable Fungicide

CEREALS – CROP GROUP 15	
CROPS	Barley, Oats
DISEASE	Barley net blotch (<i>Pyrenophora teres</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Barley, Rye
DISEASE	Barley scald (<i>Rhynchosporium secalis</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Wheat, Barley, Rye, Oats, Triticale
DISEASE	Septoria leaf spot (<i>Septoria</i> sp.)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Wheat, Barley, Rye, Triticale
DISEASE	Tan spot (<i>Pyrenophora tritici-repentis</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Barley
DISEASE	Barley leaf rust (<i>Puccinia hordei</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Winter Wheat, Spring Wheat, Barley
DISEASE	Stripe rust (<i>Puccinia striiformis</i>)
PRODUCT RATE (L/ha)	0.4 - 0.5 L/ha of Propel Fungicide + 0.2 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.

CEREALS – CROP GROUP 15	
CROPS	Winter Wheat, Spring Wheat
DISEASE	Wheat leaf rust (<i>Puccinia triticina</i>)
PRODUCT RATE (L/ha)	0.4 - 0.5 L/ha of Propel Fungicide + 0.2 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
<p>Specific Use Restrictions: DO NOT make more than one application per season of this tank mixture. An additional application of Propel Fungicide can be made, if required. Refer to the Propel Fungicide label for details of the rate and timing. A total of two applications of Propel Fungicide should be applied per season either in a tank mix with QUADRIS Flowable Fungicide or alone. DO NOT apply within 45 days of harvest (45 day PHI) for grain and straw. DO NOT harvest wheat for forage. DO NOT graze or feed livestock treated forage or cut green crop for hay or silage. The tank mix of QUADRIS Flowable Fungicide and Propel Fungicide may be applied with ground or air equipment.</p> <p>GROUND APPLICATION: Apply specified rates in a minimum of 100 L of water per hectare.</p> <p>AERIAL APPLICATION: Apply specified rates in a minimum of 45 L of water per hectare.</p>	

TANK MIX - Propel FUNGICIDE with QUADRIS Flowable Fungicide

CORN	
CROPS	Field Corn, Sweet Corn, (including Seed Production), Popcorn (including Seed Production)
DISEASE	Rust (<i>Puccinia sorghi</i>) Northern Corn Leaf Blight (<i>Setosphaeria turcicum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) Eye Spot (<i>Aureobasidium zeae</i>) Grey Leaf spot (<i>Cercospora zeae-maydis</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Make first application at the first sign of disease, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development. Good spray coverage and canopy penetration are important for best results. Apply in a minimum of 45 L of water per hectare. Use the low rate of QUADRIS under low to moderate disease pressure. Use the high rate of QUADRIS only under conditions of severe disease pressure.
<p>Specific Use Restrictions: A maximum of two applications of QUADRIS Flowable Fungicide and a maximum of two applications of Propel Fungicide should be applied per season either as a tank mix or as products applied alone. DO NOT apply to field corn and field corn grown for seed after brown silk. DO NOT apply within 30 days (30 day PHI) of harvest for forage. DO NOT apply to sweet corn within 14 days of harvest (14 day PHI).</p> <p>Not all of these cereal crops have been tested for efficacy and phytotoxicity at the recommended label rates and the tank mix of QUADRIS Flowable Fungicide and Propel Fungicide should be used at the discretion of the user. The tank mix of QUADRIS Flowable Fungicide and Propel Fungicide may be applied with ground or air equipment.</p> <p>GROUND APPLICATION: Apply specified rates in a minimum of 100 L of water per hectare.</p> <p>AERIAL APPLICATION: Apply specified rates in a minimum of 45 L of water per hectare.</p>	

NOTICE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Syngenta Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Syngenta Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below.

Accordingly, the Buyer and User assumes all risks related to performance and crop tolerance arising, and agrees to hold Syngenta Canada Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

SOYBEANS GROWN FOR SEED (Minor Use)	
DISEASES	Frogeye Leaf Spot (<i>Cercospora</i> spp.) Aerial Web Blight (<i>Rhizoctonia solani</i>)
RATE/HA	500 - 760 mL
REMARKS	Apply using ground application equipment only, when disease first appears. Under severe disease pressure, make a second application 14 days after the first application. DO NOT harvest soybeans within 30 days of the last application (30 day PHI). For use only on soybeans grown for seed. Harvested soybean seed should not be used for human food or animal feed.
CANARY SEED (Minor Use)	
DISEASE	Septoria Leaf Mottle
RATE/HA	500 mL
REMARKS	For the suppression of Septoria leaf mottle. Make one application at emergence of flag leaf. Ground application only. Apply in 200 L water/ha.
DRY EDIBLE BEANS (Minor Use)	
DISEASE	Rust
RATE/HA	500 mL
REMARKS	Apply 500 mL of Propel in minimum of 200 L water/ha by ground application or in 40 - 50 L water/ha by aerial application at the first detection of disease in the field and a second application 14 to 21 days later. APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel TO DRY EDIBLE BEANS. DO NOT APPLY WITHIN 28 DAYS OF HARVEST (28 day PHI).

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that Propel Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Propel Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of Propel Fungicide or other Group 3 fungicides with different groups that control the same pathogens.

Avoid application of more than the maximum number listed in the label and consecutive sprays of Propel Fungicide or other fungicides in the same group in a season. Use tank mixtures with fungicides from a different group when such use is permitted.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.

Monitor treated fungal populations for resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

AXIAL[®], HORIZON[®], MATADOR[®], QUADRIS[®], and WARRIOR[®] are trademarks of a Syngenta Group Company. It must be noted Propel[®] is a trademark of Viterro. Viterro is a licensed user. Buctril[®], Pardner[®] and Ripcord[®] are trademarks of their respective companies.

FOR CHEMICAL EMERGENCY, SPILL, LEAK, FIRE, EXPOSURE OR ACCIDENT, CALL DAY OR NIGHT 1-800-561-8273 or CHEMTREC – DAY OR NIGHT 1 800-424-9300

1. IDENTIFICATION OF THE MATERIAL AND SUPPLIER

- 1.1 **PRODUCT IDENTIFIER:**
TRADE NAME: PROPEL®
- 1.2 **RECOMMENDED USE:** GROUP 3 FUNGICIDE
- 1.3 **DISTRIBUTED BY:**
LOVELAND PRODUCTS CANADA, INC.
789 DONNYBROOK DRIVE • DORCHESTER, ONTARIO N0L 1G5
- 1.4 **24 Hour Emergency Phone: (Chemtrec):** 1-800-424-9300 (Toll Free) - **Additional Emergency Phone** 1-800-561-8273
Loveland Technical Service: 1-800-328-4678

2. HAZARDS IDENTIFICATION

2.1 Classification of the substance or mixture

Eye Damage/Irritation	Category 2B	H320
Skin Corrosion/Irritation	Category 2	H315
Acute Toxicity – Inhalation	Category 4	H332
Acute Toxicity – Oral	Category 4	H302
Acute Toxicity – Dermal	Category 4	H312
Sensitization – Skin	Category 1B	H317
Flammable Liquids	Category 4	H227
Environmental Hazards	Category 2	H401

2.2 Label elements



- Signal word: **WARNING**
- Hazard Statement: H320 – Causes eye irritation.
H315 – Causes skin irritation.
H302 – Harmful if swallowed.
H312 – Harmful in contact with skin.
H332 – Harmful if inhaled.
H317 – May cause an allergic skin reaction.
H227 – Combustible Liquid.
H401 – Toxic to aquatic life.
- Precautionary Statement: (Prevention): P264 – Wash hands and face thoroughly after handling.
P280 – Wear protective gloves/protective clothing/eye protection/face protection.
P270 – Do not eat, drink or smoke when using this product.
P261 – Avoid breathing dust/fume/gas/mist/vapours/spray.
P271 – Use only outdoors or in a well-ventilated area.
P272 – Contaminated work clothing must not be allowed out of the workplace.
P210 – Keep away from heat/sparks/open flames/hot surfaces – No smoking.
- Precautionary Statement: (Response): P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing.
P337+P313 – If eye irritation persists: Get medical advice/attention.
P302+P352: IF ON SKIN: Wash with plenty of soap and water.
P321: Specific treatment (see Precautions and Toxicological Information on the product label).
P333+P313 – If skin irritation or rash occurs: Get medical advice/attention
P363 – Wash contaminated clothing and wash before reuse.
P301+P312: IF SWALLOWED: Call a POISON CENTRE or doctor/physician if you feel unwell.
P330 – Rinse mouth.
P362 – Take off contaminated clothing and wash it before reuse.



SAFETY DATA SHEET

SDS NUMBER: 29548-19-LPI

SDS REVISIONS: SEC. 1

DATE OF ISSUE: 10/11/19

PROPEL®
SUPERSEDES: 10/25/16

P370+P378 – In case of fire: Use foam, carbon dioxide, dry powder, halon or water fog/mist to extinguish.

Precautionary Statement: (Storage): P403 – Store in a well-ventilated place.

Precautionary Statement: (Disposal): P501 – Dispose of contents/containers in accordance with local and provincial requirements. See product label for additional information.

2.3 Other hazards

None known

3. COMPOSITION, INFORMATION ON INGREDIENTS

3.1 Substances

3.2 Mixtures

Chemical Name:	CAS No.	Concentration [%]
Propiconazole	60207-90-1	25.00
Solvent naphtha (petroleum), heavy aromatic consists of:	64742-94-5	Proprietary
Naphthalene	91-20-3	Proprietary
1,2,4-Trimethylbenzene	95-63-6	Proprietary

4. FIRST AID MEASURES

4.1 Description of First Aid Measures

General Advice: Get medical attention if symptoms occur.

- If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.
- If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.
- If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control centre or doctor for further treatment advice.
- If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

4.2 Most Important Symptoms and Effects, Acute and Delayed

Symptoms: Harmful if inhaled. Harmful if swallowed. Causes eye irritation. Causes skin irritation. Harmful if absorbed through skin.

4.3 Immediate Medical Attention and Special Treatment

Treatment: **NOTES TO PHYSICIAN:** Contains petroleum distillate. Vomiting may cause aspiration pneumonia. There is no specific antidote. Treat symptomatically

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-561-8273

Take container, label or product name with you when seeking medical attention.

5. FIRE FIGHTING MEASURES

5.1 EXTINGUISHING MEDIA:

Suitable Extinguishing Media: Use medium appropriate to surrounding fire. Dry chemical, carbon dioxide (CO₂), foam, water spray.

5.2 SPECIAL HAZARDS ARISING FROM THE SUBSTANCE OR MIXTURE:

Specific Hazards During Firefighting: Toxic, flammable fumes are released by thermal decomposition in a fire.

5.3 SPECIAL PROTECTIVE EQUIPMENT AND PRECAUTIONS FOR FIREFIGHTERS

Special Protective Equipment for Firefighters: Self-contained breathing apparatus and full protective gear should be worn in fighting large fires involving chemicals. Use water spray to keep fire exposed containers cool. Keep people away. Isolate fire and deny unnecessary entry.

6. ACCIDENTAL RELEASE MEASURES

6.1 PERSONAL PRECAUTIONS, PROTECTIVE EQUIPMENT AND EMERGENCY PROCEDURES

Personal Precautions: Avoid inhalation of vapours and spray mist and contact with skin and eyes. Ensure adequate ventilation. Wear suitable protective clothing. Keep away from heat and sources of ignition.

6.2 ENVIRONMENTAL PRECAUTIONS

Environmental Precautions: This product is toxic to fish and aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate.

6.3 METHODS AND MATERIALS FOR CONTAINMENT AND CLEAN-UP

Methods for Clean-Up: Large Spills: Stop the flow of material, if this is without risk. Dike the spilled material, where this is possible. Absorb in vermiculite, dry sand or earth and place into containers. After removal flush contaminated area thoroughly with water.
Small Spills: Wipe up with absorbent material (e.g. cloth, fleece). Clean surface thoroughly to Remove residual contamination.
Never return spills to original containers for re-use.

7. HANDLING AND STORAGE

7.1 PRECAUTIONS FOR SAFE HANDLING:

Advice on Safe Handling: Do not use, pour, spill or store near heat or open flame.
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

7.2 CONDITIONS FOR SAFE STORAGE:

Requirements for Storage Areas and Containers: Store in original container only in a well-ventilated, cool, dry, secure area. Protec from heat, sparks and flame. Do not expose containers to temperatures above 40 °C. Keep separate from other products to prevent cross contamination. Rotate stock. Clean up spilled material immediately.

National Fire Code Classification: Class IIIA Combustible Liquid.

8. EXPOSURE CONTROLS / PERSONAL PROTECTION

8.1 CONTROL PARAMETERS:

OCCUPATIONAL EXPOSURE LIMITS

U.S. Workplace Exposure Level (ACGIH) TLVs

Components	Type	Value
Naphthalene	TWA	52 mg/m ³ (Skin)
	STEL/CEIL (C)	79 mg/m ³
Trimethylbenzene, all isomers	NIOSH RELS	125 mg/m ³

U.S. Workplace Exposure Level (OSHA) PELs

Components	Type	Value
Naphthalene	TWA	50 mg/m ³

Biological limit values

ACGIH Biological Exposure Indices

Components	Value	Specimen
No listings		

8.2 EXPOSURE CONTROLS:

Engineering Measures

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

Individual Protection Measures:

- Eye / Face Protection: Goggles or shielded safety glasses are recommended.
- Skin Protection: Long sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate or viton. Socks and chemical-resistant footwear.
- Respiratory Protection: In case of inadequate ventilation or risk of inhalation of mists or vapors, use suitable respiratory equipment approved by MSHA/NIOSH. Wear respiratory protection during operations where spraying or misting occurs. If respirators are used, a program should be in place to assure compliance with national or provincial standards. Wear air supplied respiratory protection if exposure concentrations are unknown.

9. PHYSICAL AND CHEMICAL PROPERTIES

- 9.1 APPEARANCE : Liquid
- ODOUR: Aromatic.
- ODOUR THRESHOLD: No data available.
- COLOUR: Clear yellow.
- pH: 5.4 – 8.0 (1% solution @ 25 °C)
- MELTING POINT / FREEZING POINT: Freezing Point: -40 °C.
- BOILING POINT: No data available
- FLASH POINT: 143.6 °F / 62 °C (P.M.C.C).
- FLAMMABILITY (solid, gas): No data available.
- UPPER / LOWER FLAMMABILITY OR EXPLOSIVE LIMITS: Upper: 11.7% | Lower: 1.8%.
- VAPOUR PRESSURE: 4.2 x 10⁻⁷ mmHg @ 25 °C (Propiconazole Technical).
- SOLUBILITY: 0.1 g/L @ 20 °C (Propiconazole Technical).
- PARTITION CO-EFFICIENT, n-OCTANOL / WATER: Log P = 3.65 (Propiconazole Technical).
- AUTO-IGNITION TEMPERATURE: No data available.
- DECOMPOSITION TEMPERATURE: No data available.
- VISCOSITY: 5.1 cps (or mPas) @ 21 °C.
- SPECIFIC GRAVITY (Water = 1): 0.98 g/ml
- DENSITY: 0.98 kg/L

Note: These physical data are typical values based on material tested but may vary from sample to sample. Typical values should not be construed as a guaranteed analysis of any specific lot or as specification items.

10. STABILITY AND REACTIVITY

- 10.1 REACTIVITY
Stable
- 10.2 CHEMICAL STABILITY
Stable under normal temperature conditions
- 10.3 POSSIBILITY OF HAZARDOUS REACTIONS
No data available. Will not polymerize.
- 10.4 CONDITIONS TO AVOID
Heat, flames, and sparks.
- 10.5 INCOMPATIBLE MATERIALS
Nitric acid, sulfuric acid, strong oxidizing and reducing agents.
- 10.6 HAZARDOUS DECOMPOSITION PRODUCTS
Hydrogen cyanide.



SAFETY DATA SHEET

SDS NUMBER: 29548-19-LPI

SDS REVISIONS: SEC. 1

DATE OF ISSUE: 10/11/19

PROPEL®
SUPERSEDES: 10/25/16

11 TOXICOLOGICAL INFORMATION

11.3 LIKELY ROUTES OF EXPOSURE

Eyes. Skin. Ingestion. Inhalation.

LC₅₀ (rat): >1.0 mg/L (4H)

LD₅₀ Oral (rat): 2,105 mg/kg.

LD₅₀ Dermal (rat): 4,250 mg/kg

Acute Toxicity Estimates: No data available.

Eye Irritation (rabbit): Moderate irritant.

Skin Irritation (rabbit): Moderate irritant.

Inhalation: Harmful if inhaled.

Ingestion: Harmful if swallowed.

Specific Target Organ Toxicity: Liver, skin, eye (Active ingredient). CNS, liver, kidney, blood, respiratory tract, skin, eye, stomach, thyroid, urinary bladder, skin (Other ingredients).

Aspiration: No data available.

Skin Sensitization (guinea pig): Potential sensitizer

Carcinogenicity: Naphthalene has been listed as Possibly Carcinogenic to Humans (IARC-2B).

Germ Cell Mutagenicity: Not mutagenic.

Interactive Effects: None known.

12 ECOLOGICAL INFORMATION

12.3 ECOTOXICITY

The pesticide is toxic to fish and aquatic invertebrates.

Ecotoxicological Data

	Species	Test Results
Propiconazole Technical	Oncorhynchus mykiss	0.85 ppm – 96-hour LC ₅₀
	Daphnia magna	4.8 ppm – 48-hour EC ₅₀

Drift or runoff may adversely affect non-target plants.

Do not apply directly to water.

Do not contaminate water when disposing of equipment wash water.

Do not apply when weather conditions favor drift from target area.

12.2 PERSISTENCE AND DEGRADABILITY

Biodegradability: Low to moderate persistence in soil and water.

12.3 BIOACCUMULATIVE POTENTIAL

Bioaccumulation: Low.

12.4 MOBILITY IN SOIL

Low.

12.5 OTHER ADVERSE EFFECTS

Assessment: No data available.

13 DISPOSAL CONSIDERATIONS

13.1 WASTE TREATMENT METHODS

For recyclable containers: DO NOT reuse the container for any other purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site: Triple-or pressure-rinse the empty container. Dispose of rinsate in accordance with provincial requirements. Make the rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers: For disposal, the container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse the product container for any other purpose.

14 TRANSPORT INFORMATION

14.3 LAND TRANSPORT

TDG Shipping Description: NOT REGULATED

U.S. Surface Freight Classification: INSECTICIDES OR FUNGICIDES, INSECT OR ANIMAL REPELLENTS, NOI, OTHER THAN POISON (NMFC 102120; CLASS: 60)

15 REGULATORY INFORMATION

15.1 SAFETY, HEALTH AND ENVIRONMENTAL REGULATIONS

NFPA & HMIS Hazard Ratings:

NFPA		HMIS	
2	Health	0	Least
1	Flammability	1	Slight
0	Instability	2	Moderate
		3	High
		4	Severe

SARA Hazard Notification/Reporting

SARA Title III Hazard Category:	Immediate	<u>Y</u>	Fire	<u>Y</u>	Sudden Release of Pressure	<u>N</u>
	Delayed	<u>Y</u>	Reactive	<u>N</u>		

Reportable Quantity (RQ) under U.S. CERCLA: Naphthalene (CAS: 91-20-3) 100 lbs.

SARA, Title III, Section 313: Naphthalene (CAS: 91-20-3).

RCRA Waste Code: U165 (Naphthalene).

CA Proposition 65: Not applicable.

WHMIS [Canada]: Pest control products are not controlled under WHMIS. Classified D2B

Read the approved label, authorized under the Pest Control Products Act, prior to handling the pest control product.

This chemical is a pest control product registered by Health Canada Pest Management Regulatory Agency and is subject to certain labeling requirements under the Pest Control Products Act. These requirements differ from the classification criteria and hazard information required for GHS-consistent safety data sheets. Following is the hazard information required on the pest control product label.



WARNING POISON
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

16 OTHER INFORMATION

SDS STATUS: Section 1 revised.

PREPARED BY: Product Stewardship and Regulatory Affairs

REVIEWED BY: Safety, Health and Environment

PEST CONTROL PRODUCTS ACT REG. NO.: 29548

®Propel is a registered trademark

Disclaimer and Limitation of Liability: This data sheet was developed from information on the constituent materials identified herein and does not relate to the use of such materials in combination with any other material or process. No warranty is expressed or implied with respect to the completeness or ongoing accuracy of the information contained in this data sheet, and LOVELAND PRODUCTS CANADA, INC. disclaims all liability for reliance on such information. This data sheet is not a guarantee of safety. Users are responsible for ensuring that they have all current information necessary to safely use the product described by this data sheet for their specific purpose.

GROUP 3 FUNGICIDE

Propel[®]

FUNGICIDE

COMMERCIAL

Emulsifiable Concentrate for:
Broad-spectrum disease control in Wheat, Barley, Oats, Canola, Corn, Legume Vegetables including Soybeans and for the suppression of Septoria Leaf Mottle on Canaryseed.

FOR SALE FOR USE ON TIMOTHY HAY IN THE PRAIRIE PROVINCES ONLY.

GUARANTEE:

Propiconazole250 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

CAUTION



POISON

WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

REGISTRATION NO.: **29548**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L to 1000 L**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON
N1G 4Z3
Telephone: 1-877-964-3682

Label

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IF POISONING IS SUSPECTED, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote for this product. Apply symptomatic therapy. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

PRECAUTIONS

KEEP OUT OF THE REACH OF CHILDREN. DO NOT get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin sensitization reactions in certain individuals.

Wear long pants, a long sleeve shirt, shoes and socks and chemical-resistant gloves during mixing/loading, application, clean-up and repair activities. Wear protective goggles or face shield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used. **DO NOT** eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes. After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse.

DO NOT contaminate food or feed.

DO NOT allow entry into treated area for 12 hours following application (12 hr REI). See the **DIRECTIONS FOR USE** section for crop-specific restricted entry intervals.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. This pesticide is toxic to fish. DO NOT spray any body of water by direct application, drift or by cleaning and rinsing spray equipment. This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning of equipment.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application sites such as hedgerows and woodland.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL OR REFILLING:

For recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Dispose of the rinsings in accordance with provincial requirements.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

DO NOT reuse this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

For returnable containers:

DO NOT reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

It must be noted Propel[®] is a trademark of Viterra. Viterra is a licensed user.

GROUP 3 FUNGICIDE

Propel[®]

FUNGICIDE

COMMERCIAL

Emulsifiable Concentrate for:

Broad-spectrum disease control in Wheat, Barley, Oats, Canola, Corn, Legume Vegetables including Soybeans and for the suppression of Septoria Leaf Mottle on Canaryseed.

FOR SALE FOR USE ON TIMOTHY HAY IN THE PRAIRIE PROVINCES ONLY.

GUARANTEE:

Propiconazole250 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

CAUTION



POISON

WARNING
EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER

REGISTRATION NO.: **29548**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, ON
N1G 4Z3
Telephone: 1-877-964-3682

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

IF POISONING IS SUSPECTED, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

There is no specific antidote for this product. Apply symptomatic therapy. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

PRECAUTIONS

KEEP OUT OF THE REACH OF CHILDREN. DO NOT get in eyes, on skin or on clothing. Avoid breathing spray mist or vapours. Harmful if swallowed, inhaled or absorbed through the skin. Causes eye and skin irritation. This product may cause skin sensitization reactions in certain individuals.

Wear long pants, a long sleeve shirt, shoes and socks and chemical-resistant gloves during mixing/loading, application, clean-up and repair activities. Wear protective goggles or face shield when handling the concentrated product. The wearing of neoprene gloves by pilots when entering the aircraft is essential. Mechanical flagging devices must be used. **DO NOT** eat, drink or smoke during work; wash hands and face thoroughly before doing so. Immediately wash off accidental splashes of the concentrate or spray mixture from skin, clothing and out of eyes. After work, change clothing and wash entire body thoroughly. Wash contaminated working clothes separately from other laundry before reuse.

DO NOT contaminate food or feed.

DO NOT allow entry into treated area for 12 hours following application (12 hr REI). See the **DIRECTIONS FOR USE** section for crop-specific restricted entry intervals.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. This pesticide is toxic to fish. DO NOT spray any body of water by direct application, drift or by cleaning and rinsing spray equipment. This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning of equipment.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application sites such as hedgerows and woodland.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL OR REFILLING:

For recyclable containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Dispose of the rinsings in accordance with provincial requirements.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For refillable containers:

DO NOT reuse this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

For returnable containers:

DO NOT reuse this container for any other purpose. This empty container may be returned to the point of purchase (distributor/dealer) for disposal.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

DIRECTIONS FOR USE

DO NOT use in greenhouses.

A restricted entry interval (REI) of 1 day is required for workers hand-harvesting and detasseling treated corn.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

GENERAL INFORMATION

DO NOT graze animals on treated green crops within three days of application of Propel® Fungicide.

PRODUCT INFORMATION

Propel Fungicide is a broad spectrum systemic fungicide for control of a wide range of diseases on certain crops. Propel Fungicide will protect the crop from yield and quality losses due to disease. Propel Fungicide may be used in conjunction with higher seeding rates, higher fertilizer inputs, plant growth regulators and other fungicides as required.

(See "NOTE" under PRECAUTIONS)

Crop Information:

Propel Fungicide may be applied to:

- Barley, Spring
- Canary Seed
- Canola
- Corn (including Field, Seed, and Sweet)
- Legume Vegetables
- Oats
- Soybeans
- Timothy
- Wheat, Spring and Winter (including Hard Red, Durum, Canada Prairie, Soft White)

Diseases Controlled:

Barley, Spring:	Net Blotch; Spot Blotch; Scald; Powdery Mildew; Septoria Leaf Spot; Leaf Rust; Stem Rust
Canary Seed:	Septoria Leaf Mottle
Canola:	Blackleg
Corn:	Rusts; Northern Corn Leaf Blight; Southern Corn Leaf Blight; Helminthosporium Leaf Spot; Eye Spot; Grey Leaf Spot
Legume Vegetables including Soybeans:	Rust (<i>Phakopsora</i> spp.); Cercospora Leaf Spot (<i>Cercospora</i> spp.); Powdery mildew (<i>Microsphaera</i> spp., <i>Erysiphe</i> spp.)
Oats:	Septoria Leaf Blotch; Crown Rust
Soybeans grown for seed:	Frogeye Leaf Spot (<i>Cercospora</i> spp.), Aerial Web Blight (<i>Rhizoctonia solani</i>)
Timothy:	Purple eyespot (<i>Cladosporium phlei</i>)
Wheat, Spring and Winter: (including Hard Red, Durum, Canada Prairie, Soft White)	Septoria Leaf Spot; Septoria Glume Blotch; Powdery Mildew; Leaf and Stem Rust; Tan Spot; Stripe Rust

Factors Affecting Propel Fungicide Performance:

Propel Fungicide should be applied as a preventative disease control measure. Established diseases are more difficult to control and may have already reduced crop vigour. If rainfall occurs within one hour of application, reapplication is necessary.

Field Sprayer Application:

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial Application:

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor span. High humidity and low temperatures (10 - 20°C) allow for a better deposition of spray droplets.

Sprayer and Application Information:

The performance of this product depends on correct application. Follow the guidelines given below for optimal application of Propel Fungicide.

Sprayer Information:

	GROUND APPLICATION	AERIAL APPLICATION
Spray Volume:	minimum 200 L water/ha	40 - 50 L water/ha
Spray Pressure:	200 - 300 kPa	100 - 200 kPa
Nozzle Type:	110° Flat Fan (XR11004, 4110-20)	Flat Fan (6510-6515) or Hollow Cone (D8-45)
Droplet Size:	Medium Spray (300 - 400 microns VMD)	Medium Spray (350 - 400 microns VMD)
Ground Speed:	10 km/h	
Nozzle Angle:	90° (straight down)	90° (straight down)
Boom Height:	40 - 50 cm above the crop canopy	2 - 3 m above the crop canopy

Ground Application - Mixing and Spraying Instructions

- 1) Spray equipment should be thoroughly flushed with clean water before mixing Propel Fungicide.
- 2) Fill spray tank 1/2 full with clean water. Engage gentle agitation.
- 3) Add the required amount of Propel Fungicide and agitate thoroughly.
- 4) Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tankmix partner.
- 5) Complete filling the spray tank with water, maintaining agitation during mixing and spraying operations.
- 6) Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Aerial Application - Mixing and Spraying Instructions

- 1) Spray equipment should be thoroughly flushed with clean water before mixing Propel Fungicide.
- 2) Fill premix tank 1/2 full with clean water. Engage gentle agitation.
- 3) Add the required amount of Propel Fungicide and agitate thoroughly.
- 4) Continue filling the tank with water until the tank is 9/10 full and, if applicable, add the required amount of tankmix partner.
- 5) Complete filling the premix tank with water.

- 6) Maintain gentle agitation during mixing.
- 7) Transfer the premix contents into the aircraft spray tank.
- 8) Maintain sufficient agitation during the mixing and spraying operation to ensure Propel Fungicide remains in suspension.
- 9) Use nozzle screens no finer than 50 mesh. Keep by-pass line on or near the bottom of the tank to minimize foaming.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for a specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

AERIAL APPLICATION USE PRECAUTIONS:

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift; therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as a shelter-belt) or aquatic habitat.

AERIAL APPLICATION OPERATOR PRECAUTIONS:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the label recommendations on the ground application section. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Buffer zones:

Use of the following spray methods or equipment. **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				Terrestrial habitat
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer*	Beans, peas, soybeans, chickpeas, corn, wheat, oats, canary seed, canola, barley, rye, triticale		1	0	1	1	1
Aerial	Beans, corn, oats, wheat, barley, triticale (seed prod.)	Fixed wing	1	0	3	1	20
		Rotary wing	1	0	1	1	20

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

For tank mixes, consult the labels of tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

PRODUCT SPECIFIC PRECAUTIONS:

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-87-SYNGENTA (1-877-964-3682) or obtain technical advice from the distributor or your provincial agricultural representative.

















Fertilizer - Mixing and Spraying Instructions

If desired, small amounts of nitrogen may be applied with Propel Fungicide. The appropriate amount of urea can be dissolved in water and added to the spray tank before adding Propel Fungicide. The rate of actual nitrogen must not exceed 10 kg/ha.

CAUTION: Excessive nitrogen concentrations may injure the crop.

NOTE: DO NOT add nitrogen when tank-mixing Propel Fungicide with a herbicide.

**WHEAT, BARLEY, OATS
CEREAL GROWTH STAGES***

							
3 leaves unfolded 13	Main shoot and tiller 21	Main shoot and 4 tillers 24	Leaf sheath lengthening 28	Pseudo stem anthesis 30	1st node detectable 31	2nd node detectable 32	Flag leaf just visible 37
							
Ligule of last leaf visible 39	Boots swollen 45	Ear just visible 51	Ear 1/2 emerged 55	Ear fully emerged 59	Flowering over at base of ear 69	Flowering over (anthesis) watery ripe 71	Grain hard 81

WHEAT, BARLEY, OATS	
INSTRUCTIONS FOR USE:	
Apply Propel Fungicide at the very early stages of disease. This could occur anytime during tillering or stem elongation. Typically, an application from the beginning of stem elongation up to flag leaf emergence is required.	
Propel Fungicide lasts about three weeks in the plant. If conditions favourable to disease continue after this length of time, another application will be necessary to maintain control. The second spray is usually applied at the time of head emergence. In most cases, this second application is essential to maintain control of the <i>Septoria</i> disease complex.	
LAST APPLICATION MUST BE MADE PRIOR TO 45 DAYS BEFORE HARVEST (45 day PHI).	
CROP/DISEASES	<u>WHEAT: Septoria Leaf Spot Tan Spot</u> <u>SPRING BARLEY: Net Blotch</u>
RATE/HA	250 - 500 mL
EARLY Application	At G.S. 12-23 (as early as the two leaf stage). For early season disease suppression, use the lower rate for suppression under normal field conditions. Use the higher rate for control if there is a history of high disease pressures in the field and/or field conditions favour disease development.
LATER Application	At the first sign of disease (G.S. 29-37) or before head is half emerged (G.S. 49-55). Apply only the high rate on any application from G.S. 29-55.
MAXIMUM NUMBER OF APPLICATIONS PER SEASON	2
CROP/DISEASES	<u>WHEAT: Septoria Leaf Spot and Glume Blotch Powdery Mildew Leaf and Stem Rust Tan Spot Stripe Rust</u> <u>SPRING BARLEY: Net Blotch Spot Blotch Scald Powdery Mildew Septoria Leaf Spot Leaf and Stem Rust</u> <u>OATS: Septoria Leaf Blotch Crown Rust</u>
RATE/HA	500 mL
EARLY Application	At the first sign of disease, usually at the beginning of stem elongation (G.S. 29-37).
LATER Application	Before head is half emerged (G.S. 49-55).
MAXIMUM NUMBER OF APPLICATIONS PER SEASON	2

HERBICIDE TANK-MIXING - WHEAT & BARLEY:

Propel Fungicide can be tank-mixed with ONLY ONE of these herbicides:

2,4-D Amine MCPA Amine
 Buctril® M Pardner®
 HORIZON® 240EC Herbicide (Wheat Only)
 AXIAL® 100EC Herbicide (Spring Wheat and Spring Barley Only) (FOR USE ONLY IN THE PRAIRIE AND INTERIOR OF BRITISH COLUMBIA)

Tank-mixing Precautions:

1. DO NOT tank-mix Propel Fungicide with herbicides for application onto Oats.
2. Weeds and crops must be at the correct stage of growth as specified in both the Propel Fungicide label and the tank mix partner label.
3. 2,4-D and MCPA formulations may be applied either by ground application or aerial

application; tank-mixtures of Propel Fungicide and Buctril M, Pardner, or AXIAL 100EC Herbicide can only be applied by ground application.

4. For fields treated with AXIAL 100EC Herbicide, no crop may be seeded until the following year. There are no crop rotation limitations the year following application of Axial 100EC Herbicide.
5. Tank mixing with the herbicide AXIAL 100EC will slightly decrease Propel efficacy.
6. Consult the label of the herbicide partner for a list of weeds controlled, directions for use and precautions.
7. When tank mixing, adhere to the most restrictive label limitations and precautions.
8. Compatibility should always be confirmed by premixing small proportional quantities of water, Propel Fungicide, and the tank-mix partner in advance.

NOTE:

- DO NOT graze animals on treated green crops within three days of application of Propel Fungicide.
- DO NOT feed straw from crops treated with herbicide tank mixes to livestock.

HERBICIDE TANK-MIXING – WHEAT ONLY:

FOR USE ONLY IN THE PRAIRIE AND INTERIOR OF BRITISH COLUMBIA.

Propel Fungicide can be tank-mixed with HORIZON 240EC Herbicide for disease and grassy weed control.

Tank-mixing Precautions:

1. DO NOT apply by air.
2. Consult the label of HORIZON 240EC Herbicide for a list of weeds controlled, directions for use and precautions.
3. Apply prior to emergence of the 4th tiller (herbicide timing).

TIMOTHY	
<u>INSTRUCTIONS FOR USE:</u>	
Apply Propel at the very early stages of disease. If conditions favourable to disease continue after this length of time, another application will be necessary to maintain control. DO NOT APPLY BY AIR. LAST APPLICATION MUST BE MADE PRIOR TO 14 DAYS BEFORE HARVEST (14 day PHI).	
DISEASE	Purple eyespot
RATE/HA	500 mL
EARLY Application	At the first sign of disease, usually at the beginning of flowering (G.S. 59-61).
LATER Application	Full flowering (G.S. 65-73).
Minimum Interval between Applications	14 days
Maximum Seasonal Application Rate	2 applications at 500 mL/ha (1.0 L/ha in a season).

CANOLA				
INSTRUCTIONS FOR USE:				
Propel Fungicide will control blackleg and enhance yield potential during the early stages of canola growth. The disease may reappear later in the season, but with minimal effect on yield. LAST APPLICATION MUST BE MADE PRIOR TO 60 DAYS BEFORE HARVEST (60 day PHI).				
DISEASES	RATE/HA	REMARKS/TIMING		
Blackleg	500 mL	Apply during the rosette stage: between 2 nd true leaf and bolting.		
		Seedling	Rosette	Bud (Bolted)
		Stage 1	Stage 2	Stage 3

TANK-MIXING – SEED CORN, FIELD CORN and SWEET CORN:

Propel Fungicide can be tank-mixed with ONLY ONE of the following partners:

MATADOR[®] 120EC Insecticide

WARRIOR[®] Insecticide

Ripcord[®]

Tank-mixing Precautions:

1. DO NOT APPLY THE TANK MIX WITH WARRIOR INSECTICIDE BY AIR.
2. The tank mix of Propel Fungicide + MATADOR 120EC Insecticide or Propel Fungicide + Ripcord can be applied by air and ground. Use 40 L of water per hectare when applying by air.
3. Insects and crops must be at the correct stage as specified on the Propel Fungicide as well as MATADOR 120EC, WARRIOR Insecticide and Ripcord (PCP 15738) labels. Follow the directions for use and precautions on all labels.
4. DO NOT apply within 14 days of harvest (14 day PHI) when using these tank-mixes on field and sweet corn.
5. Compatibility should always be confirmed by premixing small proportional quantities of water, Propel Fungicide, and the tank-mix partner in advance.

TANK MIXES

SEED CORN	
DISEASES	Rusts
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when rust pustules first appear. Under severe disease pressure, make a second application 14 days after. Under severe disease pressure, make a third application 14 days after. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Northern Corn Leaf Blight Southern Corn Leaf Blight Helminthosporium Leaf Spot
RATE/HA Propel Fungicide alone	250 - 500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 250 - 500 mL/ha of Propel Fungicide when disease first appears. Use the 250 mL rate if disease pressure is low. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Eye Spot Grey Leaf Spot
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when disease first appears. DO NOT apply the tank mix with WARRIOR Insecticide by air.
FIELD CORN and SWEET CORN	
DISEASES	Rusts
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when rust pustules first appear. Under severe disease pressure, make a second application 14 days after. DO NOT apply the tank mix with WARRIOR Insecticide by air.

FIELD CORN and SWEET CORN	
DISEASES	Northern Corn Leaf Blight Southern Corn Leaf Blight Helminthosporium Leaf Spot
RATE/HA Propel Fungicide alone	250 - 500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 250 - 500 mL/ha of Propel Fungicide when disease first appears. Use the 250 mL rate if disease pressure is low. DO NOT apply the tank mix with WARRIOR Insecticide by air.
DISEASES	Eye Spot Grey Leaf Spot
RATE/HA Propel Fungicide alone	500 mL
RATE/HA Tank Mix partner	+ 83 mL MATADOR Insecticide or 83 mL WARRIOR Insecticide or 175 mL Ripcord
REMARKS	Apply 500 mL/ha of Propel Fungicide when disease first appears. DO NOT apply the tank mix with WARRIOR Insecticide by air.

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS

Soybeans

Edible-podded legume vegetables - Any succulent cultivar of edible podded bean (*Phaseolus* spp.) and any succulent cultivar of edible-podded pea (*Pisum* spp.): Bean (*Phaseolus* spp.) (includes runner bean, snap bean, wax bean); bean (*Vigna* spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean); jackbean; pea (*Pisum* spp.) (includes dwarf pea, edible-podded pea, snow pea, sugar snap pea); pigeon pea; soybean (immature seed); and sword bean.

Succulent shelled pea and bean - Any succulent shelled cultivar of bean (*Phaseolus* spp.) and garden pea (*Pisum* spp.): Bean (*Phaseolus* spp.) (includes lima bean, green bean); broad bean (succulent); bean (*Vigna* spp.) (includes black-eyed pea, cowpea, southern pea); pea (*Pisum* spp.) (includes English pea, garden pea, green pea); and pigeon pea.

Dried shelled pea and bean - Any dried cultivar of bean (*Phaseolus* spp.) and dried cultivar of pea (*Pisum* spp.): Dried cultivars of bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, and sweet white lupin); (*Vigna* spp.) (includes adzuki beans); (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean, mung bean, rice bean, southern bean, urd bean); broad bean (dry); chickpea; guar; lablab bean; lentil; pea (*Pisum* spp.) (includes field pea); and pigeon pea.

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS	
DISEASE	Asian (Soybean) Rust (<i>Phakopsora pachyrhizi</i>)
CROPS	All Crop Group 6 legume vegetables listed above, and soybean
RATE/HA	500 – 756 mL
REMARKS	<p>For control of Asian Rust, apply the lower rate under conditions of low disease pressures.</p> <p>Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.</p> <p>APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).</p>
DISEASE	Powdery Mildew (<i>Microsphaera diffusa</i> , <i>Erysiphe pisi</i> , <i>E. polygoni</i>)
CROPS	All Crop Group 6 legume vegetables listed above, and soybean
RATE/HA	500 mL
REMARKS	<p>Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.</p> <p>APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).</p>
DISEASE	Cercospora Leaf Spot (<i>Cercospora kikuchi</i>)
CROPS	Soybeans only
RATE/HA	500 mL
REMARKS	<p>Make first application at the R2 to R3 developmental stage, or when there is 5% disease level in the field, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development.</p> <p>APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables).</p>

TANK MIX - Propel FUNGICIDE with QUADRIS® Flowable Fungicide

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS	
DISEASE	Asian (Soybean) Rust (<i>Phakopsora pachyrhizi</i>)
CROPS	All Crop Group 6A, B, C legume vegetables listed above, and soybeans
RATE/HA	0.5 - 0.75 L/ha of Propel Fungicide + 0.3 - 0.45 L/ha of QUADRIS Flowable Fungicide
REMARKS	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressures. A second application at 14 days interval may be needed if conditions persist. It is important to protect the developing pod of soybean and podded legume vegetables. Good spray coverage and canopy penetration are important for best results. Apply in a minimum of 45 L of water per hectare.
Restrictions: DO NOT APPLY MORE THAN 2 APPLICATIONS OF Propel FUNGICIDE PER SEASON. DO NOT APPLY MORE THAN 2 APPLICATIONS OF QUADRIS FLOWABLE FUNGICIDE PER SEASON. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables). DO NOT make more than one application to soybean hay and dry pea hay. DO NOT apply within 14 days of harvest (14 day PHI) of soybean hay and dry pea hay. DO NOT feed dried pea vines to livestock. Not all members of the legume vegetable group have been tested for efficacy and phytotoxicity at the recommended label rates and this tank mix should be used at the discretion of the user. The tank mix of Propel Fungicide plus QUADRIS Flowable Fungicide may be applied by air or ground application equipment.	

TANK MIX - Propel FUNGICIDE with MATADOR 120EC Insecticide or WARRIOR Insecticide

LEGUME VEGETABLES (Crop Group 6) INCLUDING SOYBEANS
Propel Fungicide can be tank mixed with MATADOR 120EC Insecticide or WARRIOR Insecticide for foliar disease and insect control. Apply Propel Fungicide at a rate of 250 - 756 mL/ha in a tank mix with MATADOR 120EC Insecticide or WARRIOR Insecticide at a rate of 83 - 233 mL/ha for control of soybean aphid on soybean and at a rate of 83 mL/ha for the rest of the crops in Crop Group 6. Refer to both the Propel Fungicide, MATADOR 120EC Insecticide, or WARRIOR Insecticide labels for diseases and insects controlled, specific application instructions, and precautions. Pests and crops must be at the correct stage as specified on the Propel Fungicide, MATADOR 120EC Insecticide, or WARRIOR Insecticide labels. DO NOT apply more than 2 applications per season of this tank mix. DO NOT apply within 30 days of harvest (30 day PHI) for crop subgroup 6C (dry legume vegetables) and soybeans. DO NOT apply within 15 days of harvest (15 day PHI) for crop subgroup 6A (edible podded legume vegetables) and 6B (succulent shelled legume vegetables). DO NOT graze or harvest treated forage, straw or hay for livestock feed. Not all members of the legume vegetable group have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user. This tank mix can be applied by ground application equipment only. DO NOT APPLY THE TANK MIX WITH MATADOR 120EC Insecticide or WARRIOR Insecticide BY AIR.

TANK MIXES - Propel FUNGICIDE with QUADRIS Flowable Fungicide

CEREALS – CROP GROUP 15	
CROPS	Barley, Oats
DISEASE	Barley net blotch (<i>Pyrenophora teres</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Barley, Rye
DISEASE	Barley scald (<i>Rhynchosporium secalis</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Wheat, Barley, Rye, Oats, Triticale
DISEASE	Septoria leaf spot (<i>Septoria</i> sp.)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Wheat, Barley, Rye, Triticale
DISEASE	Tan spot (<i>Pyrenophora tritici-repentis</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Barley
DISEASE	Barley leaf rust (<i>Puccinia hordei</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
CROPS	Winter Wheat, Spring Wheat, Barley
DISEASE	Stripe rust (<i>Puccinia striiformis</i>)
PRODUCT RATE (L/ha)	0.4 - 0.5 L/ha of Propel Fungicide + 0.2 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.

CEREALS – CROP GROUP 15	
CROPS	Winter Wheat, Spring Wheat
DISEASE	Wheat leaf rust (<i>Puccinia triticina</i>)
PRODUCT RATE (L/ha)	0.4 - 0.5 L/ha of Propel Fungicide + 0.2 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). Good spray coverage and canopy penetration are important for best results.
<p>Specific Use Restrictions: DO NOT make more than one application per season of this tank mixture. An additional application of Propel Fungicide can be made, if required. Refer to the Propel Fungicide label for details of the rate and timing. A total of two applications of Propel Fungicide should be applied per season either in a tank mix with QUADRIS Flowable Fungicide or alone. DO NOT apply within 45 days of harvest (45 day PHI) for grain and straw. DO NOT harvest wheat for forage. DO NOT graze or feed livestock treated forage or cut green crop for hay or silage. The tank mix of QUADRIS Flowable Fungicide and Propel Fungicide may be applied with ground or air equipment.</p> <p>GROUND APPLICATION: Apply specified rates in a minimum of 100 L of water per hectare.</p> <p>AERIAL APPLICATION: Apply specified rates in a minimum of 45 L of water per hectare.</p>	

TANK MIX - Propel FUNGICIDE with QUADRIS Flowable Fungicide

CORN	
CROPS	Field Corn, Sweet Corn, (including Seed Production), Popcorn (including Seed Production)
DISEASE	Rust (<i>Puccinia sorghi</i>) Northern Corn Leaf Blight (<i>Setosphaeria turcicum</i>) Southern Corn Leaf Blight (<i>Cochliobolus heterostrophus</i>) Eye Spot (<i>Aureobasidium zeae</i>) Grey Leaf spot (<i>Cercospora zeae-maydis</i>)
PRODUCT RATE (L/ha)	0.5 L/ha of Propel Fungicide + 0.225 - 0.3 L/ha of QUADRIS Flowable Fungicide
APPLICATION TIMING	Make first application at the first sign of disease, followed by a second application 14 days after the first, if environmental conditions are favourable for disease development. Good spray coverage and canopy penetration are important for best results. Apply in a minimum of 45 L of water per hectare. Use the low rate of QUADRIS under low to moderate disease pressure. Use the high rate of QUADRIS only under conditions of severe disease pressure.
<p>Specific Use Restrictions: A maximum of two applications of QUADRIS Flowable Fungicide and a maximum of two applications of Propel Fungicide should be applied per season either as a tank mix or as products applied alone. DO NOT apply to field corn and field corn grown for seed after brown silk. DO NOT apply within 30 days (30 day PHI) of harvest for forage. DO NOT apply to sweet corn within 14 days of harvest (14 day PHI).</p> <p>Not all of these cereal crops have been tested for efficacy and phytotoxicity at the recommended label rates and the tank mix of QUADRIS Flowable Fungicide and Propel Fungicide should be used at the discretion of the user. The tank mix of QUADRIS Flowable Fungicide and Propel Fungicide may be applied with ground or air equipment.</p> <p>GROUND APPLICATION: Apply specified rates in a minimum of 100 L of water per hectare.</p> <p>AERIAL APPLICATION: Apply specified rates in a minimum of 45 L of water per hectare.</p>	

NOTICE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Syngenta Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Syngenta Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below.

Accordingly, the Buyer and User assumes all risks related to performance and crop tolerance arising, and agrees to hold Syngenta Canada Inc. harmless from any claims based on efficacy or phytotoxicity in connection with the uses described below.

SOYBEANS GROWN FOR SEED (Minor Use)	
DISEASES	Frogeye Leaf Spot (<i>Cercospora</i> spp.) Aerial Web Blight (<i>Rhizoctonia solani</i>)
RATE/HA	500 - 760 mL
REMARKS	Apply using ground application equipment only, when disease first appears. Under severe disease pressure, make a second application 14 days after the first application. DO NOT harvest soybeans within 30 days of the last application (30 day PHI). For use only on soybeans grown for seed. Harvested soybean seed should not be used for human food or animal feed.
CANARY SEED (Minor Use)	
DISEASE	Septoria Leaf Mottle
RATE/HA	500 mL
REMARKS	For the suppression of Septoria leaf mottle. Make one application at emergence of flag leaf. Ground application only. Apply in 200 L water/ha.
DRY EDIBLE BEANS (Minor Use)	
DISEASE	Rust
RATE/HA	500 mL
REMARKS	Apply 500 mL of Propel in minimum of 200 L water/ha by ground application or in 40 - 50 L water/ha by aerial application at the first detection of disease in the field and a second application 14 to 21 days later. APPLY A MAXIMUM OF 2 APPLICATIONS PER SEASON OF Propel TO DRY EDIBLE BEANS. DO NOT APPLY WITHIN 28 DAYS OF HARVEST (28 day PHI).

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that Propel Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Propel Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of Propel Fungicide or other Group 3 fungicides with different groups that control the same pathogens.

Avoid application of more than the maximum number listed in the label and consecutive sprays of Propel Fungicide or other fungicides in the same group in a season. Use tank mixtures with fungicides from a different group when such use is permitted.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.

Monitor treated fungal populations for resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

AXIAL[®], HORIZON[®], MATADOR[®], QUADRIS[®], and WARRIOR[®] are trademarks of a Syngenta Group Company. It must be noted Propel[®] is a trademark of Viterro. Viterro is a licensed user. Buctril[®], Pardner[®] and Ripcord[®] are trademarks of their respective companies.